

## The invariant measure for $SU(N)$

Michael Creutz, Brookhaven Lab

The invariant measure for integration over  $SU(N)$  takes the form of a direct product of a uniform integration over the sphere  $S_{2N-1}$  and the invariant measure over  $SU(N-1)$

Parameterization of  $U \in SU(N)$

- the rows of the matrix  $U$  are three orthogonal complex vectors
- factor  $U$  into two pieces

$$U = \begin{pmatrix} 1 & \cdots \\ \vdots & g_{N-1} \end{pmatrix} g_s(\vec{z})$$

- $g_{N-1} \in SU(N-1)$
- $g_s \in SU(N)$  is a standard form with given top row  $\vec{z}$   $g_x(\vec{z}) = \begin{pmatrix} \vec{z} \\ \vdots \end{pmatrix}$
- $\vec{z}$  is a complex unit  $N$ -vector,  $|\vec{z}|^2 = 1$

$\vec{z}$  defines a sphere  $S_{2N-1}$

- parameterize

$$\vec{z} = \left( c_1 p_1, s_1 c_2 p_2, s_1 s_2 c_3 p_3, \dots, \prod_i^{N-2} s_i c_{N-1} p_{N-1}, \prod_i^{N-1} s_i p_N \right)$$

- $c_i = \cos(\theta_i)$ ,  $s_i = \sin(\theta_i)$ ,  $p_i = e^{i\phi_i}$
- $2N - 1$  parameters:  $\theta_1, \dots, \theta_{N-1}$ ,  $\phi_1, \dots, \phi_N$
- $0 \leq \theta_i < \pi/2$        $0 \leq \phi_i < 2\pi$

Uniform measure on the sphere

$$dS_{2N-1} = \frac{(2N-2)(2N-4)\dots 2}{(2\pi)^N} (d\theta)(d\phi) \prod_{i=1}^{N-1} c_i s_i^{2N-2i-1}$$

Parameterize the standard form  $g_s(\vec{z})$

$$g_s = \begin{pmatrix} 1 & 0 & \dots & \\ 0 & 1 & \dots & \\ \vdots & \vdots & \ddots & \\ & & & \prod_i p_i^* \end{pmatrix} \begin{pmatrix} c_1 & s_1 & 0 & \dots \\ -s_1 & c_1 & 0 & \dots \\ 0 & 0 & 1 & \dots \\ \vdots & \vdots & \vdots & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots \\ 0 & c_2 & s_2 & \dots \\ 0 & -s_2 & c_2 & \dots \\ \vdots & \vdots & \vdots & 1 \end{pmatrix} \\ \dots \begin{pmatrix} 1 & \vdots & \vdots & \vdots \\ \dots & 1 & 0 & 0 \\ \dots & 0 & c_{N-1} & s_{N-1} \\ \dots & 0 & -s_{N-1} & c_{N-1} \end{pmatrix} \begin{pmatrix} p_1 & 0 & \dots \\ 0 & p_2 & \dots \\ \vdots & \vdots & \ddots \\ & & & p_N \end{pmatrix}$$

This parameterization covers the group

- $g_{N-1}$ :  $(N - 1)^2 - 1$  parameters
- $S_{2N-1}$ :  $2N - 1$  parameters
- total:  $N^2 - 1$  parameters of  $SU(N)$

Measure for  $SU(N)$

- $dg_N = f(\vec{z}, g_{N-1}) dS_{2N-1} dg_{N-1}$
- weight factor  $f(\vec{z}, g_{N-1})$  to be determined

Left invariance over the  $SU(N - 1)$

- $f(\vec{z}, g_{N-1})$  cannot depend on  $g_{N-1}$

Right invariance on  $SU(2)$  subgroups

- can implement arbitrary rotations on  $\vec{z}$
- measure does not depend on  $\vec{z}$

$$dg_N = dS_{2N-1} dg_{N-1}$$

Measure uniform over both  $S_{2N-1}$  and  $g_{N-1}$

## Example: $SU(3)$

$$g = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & p_1^* p_2^* p_3^* p_4^* p_5^* \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_3 & s_3 \\ 0 & -s_3 & c_3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & p_4 & 0 \\ 0 & 0 & p_5 \end{pmatrix}$$

$$\begin{pmatrix} c_1 & s_1 & 0 \\ -s_1 & c_1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_2 & s_2 \\ 0 & -s_2 & c_2 \end{pmatrix} \begin{pmatrix} p_1 & 0 & 0 \\ 0 & p_2 & 0 \\ 0 & 0 & p_3 \end{pmatrix}$$

Five phases and three angles

$$dg = \frac{3}{8\pi^5} c_1 c_2 c_3 s_1^2 s_2 s_3 (d\theta)(d\phi)$$

$$0 \leq \theta_i < \pi \quad 0 \leq \phi_i < 2\pi$$

## Periodicity and $\Pi_{2N-1}(SU(N))$

Map  $S_{2N-1}$  into the group in a smooth but non-contractable way

Try mapping the  $S_{2N-1}$  into the top row

- parameterization of  $g_s$  singular at the poles when  $s_i = 0$
- resolve recursively by defining a new  $\tilde{g}_s =$

$$\begin{pmatrix} p_1 & 0 & 0 & \dots \\ 0 & 1 & 0 & \dots \\ 0 & 0 & 1 & \dots \\ \vdots & \vdots & \vdots & p_1^* \end{pmatrix} \begin{pmatrix} 1 & \dots \\ \vdots & g_{N-1}^\dagger \end{pmatrix} \begin{pmatrix} c_1 & s_1 & 0 & \dots \\ -s_1 & c_1 & 0 & \dots \\ 0 & 0 & 1 & \dots \\ \vdots & \vdots & \vdots & 1 \end{pmatrix} \begin{pmatrix} 1 & \dots \\ \vdots & g_{N-1} \end{pmatrix}$$

- $g_{N-1}$  submatrix is in  $SU(N-1)$  and contains the remaining angles

At the north pole with  $\theta_1 = 0$

- the second factor cancels the last
- things are smooth

But, still singular at the south pole

- cut out that pole
- keep going in  $\theta_1$  beyond  $\pi/2$
- at  $\theta_1 = \pi$  things are still singular
- finally at  $\theta_1 = 2\pi$  get a non singular point

For a nontrivial mapping of  $S_{2N-1}$  into  $SU(N)$

- take  $\theta_1 = \tilde{\theta}_1/4$
- $\tilde{\theta}_1$  parameterizes the  $S_{2N-1}$  top row
- for  $\Pi_{2N-1}(SU(N))$  we cover the range of  $\tilde{\theta}_1$  4 times

Recurring to the lower groups

- additional factors of 4 until  $SU(2)$
- a  $SU(2)$  matrix is entirely determined by its top row

To map an  $S_{2N-1}$  nontrivially into  $SU(N)$

- sweep over possible first rows  $4^{N-2}$  times

$$\text{Bott periodicity factor} = 4^{N-2}$$