

*Work supported by the U. S. Atomic Energy Commission.

¹M. H. L. Pryce, Proc. Roy. Soc. (London) **A165**, 247 (1938).

²K. M. Case, Phys. Rev. **106**, 1316 (1957).

³N. Rosen and P. Singer, Bull. Res. Council Israel **8F**, 51 (1959).

⁴I. M. Barbour, A. Bietti, and B. F. Touschek, Nuovo Cimento **28**, 452 (1963).

⁵D. K. Sen, Nuovo Cimento **31**, 660 (1964).

⁶B. Ferretti, Nuovo Cimento **33**, 264 (1964).

⁷W. A. Perkins, Phys. Rev. **137**, B1291 (1965).

⁸B. Ferretti and I. Venturi, Nuovo Cimento **35**, 644 (1965).

⁹P. F. Smith, Nuovo Cimento **38**, 1824 (1965).

¹⁰B. Jouvét, Nuovo Cimento **38**, 951 (1965).

¹¹P. Bandyopadhyay, Nuovo Cimento **38**, 1912 (1965).

¹²V. S. Berezinskii, Zh. Eksperim. i Teor. Fiz. **51**,

1374 (1966) [Soviet Phys. JETP **24**, 927 (1967)].

¹³P. Ehrenfest and J. R. Oppenheimer, Phys. Rev. **37**, 333 (1931).

¹⁴H. L. Sahlín and J. L. Schwartz, Phys. Rev. **138**, B267 (1965).

¹⁵R. Penney, J. Math. Phys. **6**, 1031 (1965).

¹⁶See Ref. 7, Appendix A.

¹⁷L. P. Kadanoff and G. Baym, *Quantum Statistical Mechanics* (Benjamin, New York, 1962).

¹⁸F. Paschen, Ann. Physik **4**, 277 (1901).

¹⁹O. Lummers and E. Pringsheim, Verh. Deut. Physik Ges. **2**, 163 (1900).

²⁰H. Rubens and F. Kurlbaum, Ann. Physik **4**, 649 (1901).

²¹W. W. Coblentz, Natl. Bur. Std. (U.S.) Bull. **13**, 459 (1916).

²²P. Jordan, Z. Physik **93**, 464 (1935).

²³See Ref. 12, Sec. 2.

Compton Scattering and Fixed Poles in Parton Field-Theoretic Models*

Stanley J. Brodsky, Francis E. Close,[†] and J. F. Gunion

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305

(Received 12 October 1971)

We extend a class of parton models to a fully gauge-invariant theory for the full Compton amplitude. The existence of local electromagnetic interactions is shown to always give rise to a constant real part in the high-energy behavior of the amplitude $T_1(\nu, q^2)$. In the language of Reggeization this is interpreted as a fixed pole at $J=0$ in T_1 and νT_2 , with residue polynomial in the photon mass squared.

Recent inelastic electroproduction experiments (which essentially measure the imaginary part of the forward off-shell Compton amplitude) hint at a composite nature for the nucleon. This has been represented by parton models involving pointlike (possibly field-theoretic) constituents, but up to the present time these concepts have only been applied to the scaling, incoherent impulse approximation, region. Gauge invariance and the low-energy theorem place further restrictions upon such theories, and in this note we report the extension of parton field-theoretic ideas to a discussion of the full Compton amplitude. In particular we shall see that such models always give rise to a real part at high energies additional to that expected from the Regge behavior of the imaginary part. This extra real part should be identified with the "fixed pole"¹ of conventional Regge analysis. Evidence for such a fixed pole for on-shell photons has been found phenomenologically from dispersion relations.² In addition we find that the "fixed pole" appears as a constant real part, C , in T_1 independent of q^2 , and appears in νT_2 in the form

$$-Cq^2/\nu.^3$$

If the proton were as simple as the nucleus, then the high-energy behavior of the forward Compton amplitude would follow directly from the coherent impulse approximation. At $\nu=0$, the Compton amplitude on a nucleus is given by the Thomson limit⁴ $f_1(0) = -(Z^2\alpha/M_{\text{nucleus}})$ whereas at energies high compared to the binding energy, but below threshold for photoproduction of mesons, the forward amplitude is given by the coherent sum of the individual nucleon amplitudes,

$$f_1(\nu) \rightarrow -\sum_{i=1}^Z \frac{\alpha}{\omega_i} \quad (\omega_i \simeq m_i).$$

In fact, for the case of a composite proton the analogous high-energy behavior would be given by the coherent sum of "seagull" terms for the individual proton constituents (quarks, bare hadrons) and the formulas (7), (11) we give later correspond to this picture.

Field theory gives us the clearest example of a fully covariant, gauge-invariant Compton amplitude which can also incorporate the composite na-

ture of the nucleon. As an example, consider a ϕ^3 field theory in which a scalar "proton" interacts with neutral scalar particles. We calculate the form factor $F_1(q^2)$ using old-fashioned perturbation theory in an infinite-momentum frame⁵ by evaluating the matrix element of the electromagnetic current through second order from the time-ordered contributions of the Feynman diagrams 1(a)–1(c).

At $q^2=0$ we obtain

$$F_1(0) = \int_0^1 f(x) dx, \quad (1)$$

where $f(x)$ is the probability for the charged constituent to be in the one- or two-body state and to have fractional longitudinal momentum x (defined in the infinite-momentum frame).⁵ It is given by

$$f(x) = Z_2 \delta(1-x) + \frac{g^2}{(2\pi)^3} \int d^2k_\perp \frac{x(1-x)}{D^2},$$

with $D = k_\perp^2 + x\mu^2 + (1-x)M^2$, where M (μ) is the mass of the charged (uncharged) constituent and $Z_2 \equiv (1-B)^{-1} = 1 + B_{(2)} + O(g^4)$ is the familiar wave-function renormalization constant. From Figs. 1(b) and 1(c) one finds that

$$B_{(2)} = -L_{(2)} = -\frac{g^2}{(2\pi)^3} \int d^2k_\perp \int_0^1 dx \frac{x(1-x)}{D^2}$$

$$T_1(\nu, q^2) = -2 \left[Z_2 + \frac{g^2}{16\pi^3} \int d^2k_\perp \int_0^1 dx (1-x) \left(\frac{1}{D^2} - \sum_{\pm\nu} \frac{2k_i^2(1-x)}{D^2[D' + x(1-x)(q_\perp^2 - 2M\nu) - i\epsilon]} \right) \right], \quad (2)$$

where $D' = D(k'_\perp^2)$ with $\vec{k}'_\perp = \vec{k}_\perp - (1-x)\vec{q}_\perp$; k_i is the component of \vec{k}_\perp in the direction orthogonal to both \vec{p} and \vec{q}_\perp . After integrating by parts on dk_\perp^2 and taking the limit $\nu \rightarrow 0$ [for $q^2 (= -q_\perp^2) = 0$], one recovers the Thomson limit,

$$\lim_{\nu \rightarrow 0} T_1(\nu, 0) = -2 \equiv T_1^{\text{Born}}.$$

On the other hand, at large energies only the sea-gull terms [Figs. 1(d)–1(f)] survive and

$$\begin{aligned} \lim_{\nu \rightarrow \infty} T_1(\nu, q^2) &= -2 \left(Z_2 + \frac{g^2}{16\pi^3} \int d^2k_\perp \int_0^1 \frac{dx(1-x)}{D^2} \right) \\ &= T_1^{\text{Born}} \int_0^1 \frac{dx}{x} f(x). \end{aligned} \quad (3)$$

We can also compute T_2 to the same order in perturbation theory and we find the following results:

$$\begin{aligned} (i) \quad \lim_{\nu \rightarrow \infty} \nu W_2(\nu, q^2) &\left(\equiv \frac{1}{2\pi M} \text{Im} \nu T_2(\nu, q^2) \right) \\ &_{(-2M\nu/q^2) = \omega = \text{fixed}} \\ &= x f(x) \Big|_{x=1/\omega}, \end{aligned} \quad (4)$$




FIG. 1. (a)–(c) The vertex Feynman diagrams. (d)–(h) The Feynman diagrams contributing to T_1 .

and so $F_1(0) = 1$ (a consequence of the Ward identity). One can similarly sum the contributions to Compton scattering 1(d)–1(h) and obtain

$$(ii) \quad \lim_{\nu \rightarrow \infty} \nu T_2(\nu, q^2) \sim -\frac{q^2}{\nu} T_1^{\text{Born}} \int_0^1 \frac{dx}{x} f(x), \quad (5)$$

and the gauge-invariance condition $(q^2/\nu^2)T_2 + T_1 = O(q^2)$ is satisfied as $q^2 \rightarrow 0$.

Equations (1) and (3)–(5) in terms of $f(x)$ hold also for the case of spin- $\frac{1}{2}$ constituents interacting either by pseudoscalar or vector exchange, but the precise form of $f(x)$ is different for each case.⁶

These results can be generalized to all orders in perturbation theory as follows. We calculate the form factor (using the infinite-momentum frame of Ref. 5) by evaluating the matrix element of the j_0 current from the time-ordered contributions of all Feynman graphs, through arbitrary order. The contributions can be classified according to the number of intermediate constituents present at the time the currents acts [Fig. 2(a)] and the type of constituent, a , with charge $|e|\lambda_a$, upon which it acts. At $q^2=0$ we obtain⁷

$$F_1(0) = 1 = \int_0^1 dx \sum_a \lambda_a \sum_n N_a^n f_a^n(x)$$

$$\equiv \int_0^1 dx \sum_a \lambda_a f_a(x), \quad (6)$$

where N_a^n is the multiplicity of parton type a in the n -constituent state. Provided $Z_2 \neq 0$, $f_a^n(x)$ retains its interpretation as the probability for parton type a to have fractional longitudinal momentum x and to be in the n -body state. The index a runs over both the partons and antipartons present. If $Z_2 = 0$, then $f_a^n(x)$ may lose its interpretation as a probability; $\sum_n \int f_a^n(x) dx$ can be infinite. The generalized seagull, or Z graph, contribution [Fig. 2(b)] yields a constant term in T_1 at high energy,

$$\begin{aligned} T_1^{\text{FP}} &= T_1^{\text{Born}} \sum_a \lambda_a^2 \int_0^1 \frac{f_a(x)}{x} dx \\ &= -\frac{\nu^2}{q^2} T_2^{\text{FP}}, \end{aligned} \quad (7)$$

analogous to the nuclear-physics result. (Here "FP" means "fixed pole.") In the scaling limit we can identify

$$\nu W_2(x) = x \sum_a \lambda_a^2 f_a(x) \equiv x f(x) \quad [\text{Fig. 2(c)}]. \quad (8)$$

The question of convergence of Eq. (7) is very important and must be considered carefully. On the one hand, it is possible that $f(x)$ is well behaved and vanishes as $x \rightarrow 0$. In this case Regge behavior in the structure function νW_2 cannot result from the parton distribution. Equation (7) then gives the exact parton contribution to the




FIG. 2. Generalized time-ordered diagram for the (a) form factor; (b) "seagull"; (c) νW_2 . The matrix elements are proportional to $\sum_a \lambda_a N_a$, $\sum (\lambda_a^2/x_a) N_a$, $\sum \lambda_a^2 N_a x_a \delta(x_a - Q^2/2M\nu)$, respectively. N_a is the number operator for parton a .

constant term in T_1 and $-(\nu^2/q^2)T_2$. On the other hand, the parton distribution for small x (i.e., wee partons) may give rise to leading Regge behavior as discussed in Refs. 8 and 9. In such a case $f(x) \sim \sum_{\alpha>0} x^{-\alpha} \gamma_\alpha$ for $x \sim 0$ and the integral in Eq. (7) diverges. [The integral for $F_1(0)$ does not diverge because of the cancellation of the Pomeron $\alpha=1$ contribution, and in general all $C=+$ exchanges, of a given parton with the analogous contribution of its antiparton.] One can see from Eq. (2) that the nonseagull terms play an important role in removing the apparent divergence at $x=0$. In fact, despite the presence of Regge terms, the fixed pole - constant real part - survives in a slightly altered form. To see this we now turn to the covariant, nonperturbative parton model developed in Ref. 9, which allows one to incorporate leading Regge behavior in a natural way.

We restrict ourselves to spinless partons for which the required distribution function is

$$f_a(x) = \frac{Ax}{x-1} \int ds d^2K \text{Im} T_R^a(s, \mu_s^2), \quad (9)$$

$\mu_s^2 = x(s - K^2)/(x-1) + xM^2 + K^2$, where K is space-like, two-dimensional, and orthogonal to p and q ; and μ_s^2 is the invariant four-momentum squared of the interacting parton. The integral over s is over the right-hand cut of the forward parton-proton scattering amplitude, $\text{Im} T_R^a(s, \mu^2)$ [Fig. 3(a)] (which includes the propagators of the partons and is assumed to vanish as $\mu^2 \rightarrow \infty$). That $f(x)$ is related to the forward parton-proton scattering amplitude is already apparent in the perturbation-theory approach.¹⁰ One notices that for small x ($\mu_s^2 \equiv -z + K^2$)




FIG. 3. (a) The parton-proton scattering amplitude. (b) The generalized seagull contribution. (c) The freely propagating parton graphs. [The self-energy modifications included in $\rho_a(m^2)$, Eq. (11), are not drawn.] (d) The fully connected diagram.

$$f^a(x) \rightarrow Ax^{-\alpha} \int_0^\infty dz \int d^2K z^\alpha \beta_\alpha^a(-z+K^2) = x^{-\alpha} \gamma_\alpha^a$$

if

$$\text{Im}T_R^a(s, \mu^2) \sim s^\alpha \beta_\alpha^a(\mu^2).$$

One can calculate in a similar fashion the contribution for the antiparton amplitude. Note the assumption that the parton-nucleon amplitude has normal Regge behavior. The resulting expressions for $F_1(0)$, the seagull term, and νW_2 are formally

$$A \int_0^1 dx \int dm^2 \rho_a(m^2) \int d^2K K^2 \int ds \left(\frac{1}{\mu_s^2 + 2xM\nu - m^2} + \frac{1}{\mu_s^2 - 2xM\nu - m^2} \right) \left(\frac{1}{x-1} \text{Im}T_R^a(s, \mu_s^2) \right), \quad (10)$$

where $\rho_a(m^2)$ is the spectral function for the propagator of parton a . As in the second-order perturbation-theory case, a partial-integration identity on $\text{Im}T_R^a$ must be obeyed (in order that the low-energy theorem be satisfied). One then finds that for large ν the sum of these terms and the seagull gives a constant real part to T_1 of the form (for all q^2 and ν)

$$T_1^{\text{FP}} = \sum_a \lambda_a^2 \left(\int_0^1 \frac{\tilde{f}^a(x)}{x} dx - \sum_{\alpha>0} \frac{1}{\alpha} \gamma_\alpha^a \right) \times T_1^{\text{Born}}, \quad (11)$$

where the \tilde{f}^a are defined by

$$\tilde{f}^a(x) = f^a(x) - \sum_{\alpha>0} \gamma_\alpha^a x^{-\alpha}.$$

The remaining contributions from the above diagrams, and also the total contribution of diagram 3(d), yield conventional leading Regge behavior with the expected phases. As before we also find a corresponding real part in $\nu T_2 = -(q^2/\nu) T_1^{\text{FP}}$.

Thus a fixed pole arises in the Compton amplitude due to local photon interactions even though the parton-proton amplitude has no such term.

The q^2 -independent value of the fixed pole in Eq. (11) is precisely the finite-energy sum-rule (FESR) result which one obtains in the scaling limit. Explicitly the FESR (in the scaling limit) is

$$\int_0^1 \frac{\tilde{f}(x)}{x} dx - \sum_{\alpha>0} \frac{1}{\alpha} \gamma_\alpha \equiv \int_0^\infty \tilde{F}_2(\omega) d\omega \stackrel{(\text{FESR})}{=} \frac{\nu^2}{2q^2} T_2^{\text{FP}},$$

$$\tilde{f} = \sum_a \lambda_a^2 \tilde{f}_a, \quad \gamma_a \equiv \sum_a \lambda_a^2 \gamma_\alpha^a,$$

which is precisely the result we obtain for all q^2 , $\nu \rightarrow \infty$.¹²

Thus we find that fixed-pole - real-part contributions are always associated with the existence of seagull or corresponding Z -graph couplings to the charged constituents of the target. Apart from the term $\sum_\alpha (1/\alpha) \gamma_\alpha$ which results from the conventional Reggeization procedure, only $\tilde{f}(x)$, that part of the distribution function which does not con-

tain the leading (i.e., divergent) x behavior $\sum_\alpha x^{-\alpha} \gamma_\alpha$ contributes. The remaining part is absorbed into the normal leading Regge behavior of the full Compton amplitude. The distribution function $\tilde{f}(x)$, which vanishes as $x \rightarrow 0$, may be associated with short-range terms in the space-time structure of the current correlation function, as shown by suri and Yennie.⁶ We note that the seagull (or corresponding Z graph) contribution to the real part of the general Compton amplitude $T_{\mu\nu}(q_1^2, q_2^2, s, t)$ is independent of the photon masses and depends only upon $t = (q_1 - q_2)^2$ in the form

$$g_{\mu\nu} \sum_a \lambda_a^2 \int_0^1 \frac{f_a(x, t)}{x} dx$$

and has dependence on t similar to that of the elastic form factor,¹³

$$\sum_a \lambda_a \int_0^1 f_a(x, t) dx = F_1(t).$$

As before leading Regge terms must be subtracted from the seagull contribution to obtain the fixed pole. This form can be tested in both nonforward (elastic or inelastic) Compton scattering and photo-production of lepton pairs, since the leading Regge contributions are expected to disappear much more rapidly than the fixed-pole contribution as t grows. Our results also have interesting implications for the processes $\gamma + \gamma \rightarrow X$ which are accessible in $ee \rightarrow eeX$ measurements.¹⁰

In the case of a simple three-quark model of the nucleon with the same distribution function for \mathcal{P} and \mathcal{N} quarks, then $T_1^{\text{FP}}(n) = \frac{2}{3} T_1^{\text{FP}}(p)$.¹⁴ In general a composite theory of the neutron with charged constituents leads to a nonzero fixed-pole contribution; accordingly, direct measurements of the real part of the nucleon Compton amplitude from Bethe-Heitler interference experiments for both proton and deuteron targets will be very interesting.

We thank our colleagues at SLAC for interesting and helpful discussions.

*Work supported in part by the U. S. Atomic Energy Commission.

†NATO Fellow 1970–1972.

¹The forward Compton amplitude is written $\epsilon^\mu \epsilon^\nu T_{\mu\nu}$, where

$$T_{\mu\nu} = - \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) T_1(q^2, \nu) + \frac{1}{M^2} \left(P_\mu - \frac{P \cdot q}{q^2} q_\mu \right) \left(P_\nu - \frac{P \cdot q}{q^2} q_\nu \right) T_2(q^2, \nu).$$

Both $T_{1,2}$ are crossing-symmetric so they Reggeize as

$$T_1 \sim (1 + e^{-i\pi\alpha}) \nu^\alpha + C,$$

$$T_2 \sim (1 + e^{-i\pi\alpha}) \nu^{\alpha-2} - C q^2 / \nu^2,$$

where the terms involving the real number C are possible fixed poles at $J=0$. Note that the fixed pole is defined relative to Regge terms with the correct phase.

²M. J. Creutz, S. D. Drell, and E. A. Paschos, Phys. Rev. **178**, 2300 (1969); M. Damashek and F. J. Gilman, Phys. Rev. D **1**, 1319 (1970); C. A. Dominguez, C. Ferro-Fontan, and R. Suaya, Phys. Letters **31B**, 365 (1970). In the Damashek-Gilman analysis $\text{Re}T_1 \sim \nu^{1/2} + C$, where $C \sim T_1^{\text{Born}}$.

³Polynomial behavior had been previously conjectured by T. P. Cheng and Wu-Ki Tung, Phys. Rev. Letters **24**, 851 (1970), and J. M. Cornwall, D. Corrigan, and R. Norton, *ibid.* **24**, 1141 (1970). The constraints of such behavior for the electroproduction data have been discussed by F. E. Close and J. F. Gunion, Phys. Rev. D **4**, 742 (1971); **4**, 1576 (1971). It should be noted that polynomial residue plus finiteness as $q^2 \rightarrow \infty$ yields the form described in the text.

⁴The normalization used is $f_1(\nu) = (\alpha/2M) T_1(\nu, 0)$.

⁵The frame chosen is such that

$$p_\mu = (P + M^2/2P, \vec{0}, P),$$

$$q_\mu = \left(\frac{M\nu}{2P}, \vec{q}_\perp, \frac{-M\nu}{2P} \right),$$

and internal integration (on-mass-shell) four-vectors are chosen in the form

$$k_\mu = \left(|1-x|P + \frac{k_\perp^2 + M^2}{2|1-x|P}, -\vec{k}_\perp, (1-x)P \right).$$

The only kinematic conditions imposed are $Q^2 = -q^2 > 0$, $p^2 = M^2$, $p \cdot q = M\nu$, and terms of order M^2/P^2 , $M\nu/P^2$ are neglected as $P \rightarrow \infty$. Note that only one time ordering of the Feynman diagrams 1(a)–1(h) survives in ϕ^3 theory as $P \rightarrow \infty$. See S. Weinberg, Phys. Rev. **150**, 1313 (1966). Many of the techniques used in this paper were developed by Drell, Levy, and Yan, Refs. 7 and 13.

⁶For spin- $\frac{1}{2}$ particles, the “seagull” arises from a Z graph in the infinite-momentum frame. [See Fig. 2(b).] Covariant regularization must be used in these theories

to guarantee convergence, proper gauge invariance, and low-energy theorems. Note also that to any finite order in the perturbation theory for any regularized theory $f(x)$ vanishes as $x \rightarrow 0$, and T_1 and T_2 do not have leading Regge behavior. This exhibits the behavior of the short-range component in νW_2 as discussed by A. suri and D. R. Yennie, SLAC Report No. SLAC-PUB-954 (unpublished). We thank Dr. suri and Dr. Yennie for helpful conversations.

⁷In this sum of n -particle states, the $n=1$ state is presumed absent in order to ensure that the elastic form factor vanishes as $q^2 \rightarrow \infty$. This is equivalent to assuming the dynamical requirement $Z_2 \rightarrow 0$ [i.e., dropping Born terms in the perturbation-theory calculation – see S. Drell, D. Levy, and T. M. Yan, Phys. Rev. **187**, 2159 (1969); Phys. Rev. D **1**, 1035 (1970)].

⁸J. D. Bjorken, Phys. Rev. **179**, 1547 (1969); R. P. Feynman, Phys. Rev. Letters **23**, 1415 (1969); J. D. Bjorken and E. A. Paschos, Phys. Rev. **185**, 1975 (1969); J. Kuti and V. F. Weisskopf, Phys. Rev. D **4**, 3418 (1971).

⁹P. V. Landshoff, J. C. Polkinghorne, and R. D. Short, Nucl. Phys. **B28**, 222 (1971). In the normalization of this reference $A = -[2/(2\pi)^3] Z_2^2$, where Z_2^2 is the wavefunction renormalization constant for parton a . T is then a fully renormalized amplitude.

¹⁰See S. J. Brodsky, F. E. Close, and J. F. Gunion (unpublished), in which an exhaustive account of the results of this note will be presented.

¹¹For $q^2 \neq 0$, the divergence at $x=0$ [originating from the fact that $f_a(x) \sim x^{-\alpha} \gamma_\alpha^a$ as $x \rightarrow 0$] is canceled by a subtraction term which results when a subtracted dispersion representation is used for the leading Regge terms of the parton-proton $T^{(4)}$ amplitude. This results in the replacement of $f(x)$ by $\tilde{f}(x)$ and gives rise to the $-(1/\alpha)\gamma_\alpha$ term in Eq. (11). At $q^2=0$ the effects of the subtraction term disappear.

¹²This gives a physical realization of the result obtained by Cornwall, Corrigan, and Norton, Ref. 3.

¹³S. D. Drell and T. M. Yan, Phys. Rev. Letters **24**, 181 (1970).

¹⁴In such a simple three-quark model $T_1^{\text{FP}} \sim 3T_1^{\text{Born}}$ (taking $\langle 1/x \rangle \cong 3$) in an energy region where the impulse approximation is relevant. This does not agree with the FESR dispersion analyses in Ref. 2, but it is quite possible that the assumptions on asymptotic behavior in these analyses are incorrect if a threshold for quark production exists which has yet to be reached. If this were the case then direct measurements of the real part of the Compton amplitude should show a slow transition to the ultimate asymptotic result. This of course would make the impulse approximation used to explain scaling hard to understand. Further discussion will be given elsewhere.